

Study on phase transition critical phenomena in group movement based on eigen microstate theory

Weihang Wang

School of System, Beijing Normal University, Beijing, China, 100878

wwh.flame1837@gmail.com

Abstract. Group motion exists widely in nature. Physicists use abstract models, such as the Vicsek model, to simplify group motion to obtain universal laws. However, the order parameters and phase transition types that describe the characteristics of the Vicsek model system are still controversial. For example, how to characterize the order parameters and types of phase transitions that characterize the system. This paper starts with the eigen microstate method and studies the Vicsek model. First, the continuous phase transition in the system is deeply studied through scaling analysis, and two critical exponents about the noise are obtained. Second, considering the density effect, similar to the noise study, three eigen microstates in the Vicsek model are revealed through scaling analysis, showing discontinuous and continuous phase transitions. Moreover, it is proved that in the actual infinite system, continuous phase transition and discontinuous phase transition still exist. Finally, the experimental data under different noises and densities are obtained through a large number of numerical simulations, and the phase diagram of the Vicsek model is depicted.

Keywords: group motion, eigen microscopic state, phase transition critical phenomenon, noise, density, phase diagram

1. Introduction

Group movement is an orderly movement that occurs spontaneously in a system composed of many self-driven units [1-3]. These phenomena are studied through experiments, simulations [4], and theory [5, 6]. One of the most famous models is the Vicsek model introduced by Tamás Vicsek et al. in 1995 [7].

In previous work, the average velocity of all particle motions was used as the order parameter of the Vicsek [7-9] model. Tamás Vicsek and others believe that the type of phase transition that occurs in the Vicsek model is a continuous phase transition. However, the continuity of the Vicsek model phase transition has been questioned by Chaté et al. [8, 9]. Chaté et al. believe that in real systems, the Vicsek model has this discontinuity characteristic. Therefore, the continuity of phase transitions in the Vicsek model is not exactly defined.

Huepe et al. [10] questioned the use of velocity as the order parameter of the Vicsek model. They discussed the number density of particles that act on particles as the order parameter of the Vicsek model, and considered that the phase transition type of the Vicsek model is continuous phase transition.

Due to disputes over the order parameters and phase transition types of the Vicsek model, continuous phase transitions and discontinuous phase transitions appear in different regions in the phase diagram of the Vicsek model drawn by R. T. Wicks et al. [11].

Xu Li et al. applied the eigen microstate method to the study of Vicsek model and found that there are three main eigen microstates [12]. Continuous phase transition and discontinuous phase transition exist at the same time. However, Li et al. [12] only discussed the impact of noise on the Vicsek model in a small noise range, and did not discuss the impact of density on the Vicsek model, and did not calculate the specific critical exponent of the Vicsek model. It is difficult to describe the behavior of the Vicsek model in the entire phase space.

Based on this, this paper first further studies the Vicsek model under the influence of noise, and calculates the size of the specific critical exponent. Secondly, this paper discusses the behavior of the Vicsek model under the influence of density. Finally, through a large number of numerical simulations, the Vicsek model is drawn phase diagram, thereby illustrating the behavior within the phase space of the Vicsek model. And the entire phase space is divided into an ordered liquid phase, a disordered liquid phase, and a disordered gas phase.

2. Vicsek model and Eigen microstates

Tamás Vicsek et al introduced the Vicsek model in 1995 [7]. In the model, the particle is described as the direction of the particle position $r_i(t)$ and particle velocity $\Theta_i(t)$. The evolution of particles is determined by the following two equations:

$$\begin{cases} \Theta_i(t) = \langle \Theta_j(t - \Delta t) \rangle_{|r_i - r_j| < r} + \Delta\theta(t) \\ \mathbf{r}_i(t + \Delta t) = \mathbf{r}_i(t) + v_0 \Delta t \begin{pmatrix} \cos \Theta_i(t) \\ \sin \Theta_i(t) \end{pmatrix} \end{cases} \quad (1)$$

where $\langle \Theta_j(t) \rangle_{|r_i - r_j| < r}$ is the average motion direction of particles within distance r ; and $\Delta\theta(t)$ is in the range of $[-\eta/2, \eta/2]$ uniform disturbance. When the particle moves, it moves in a new direction according to the given speed v_0 .

At the initial moment, N particles will be evenly distributed in the periodic two-dimensional space of $L \times L$.

For the Vicsek model, we introduce the neighbor number density of particle i at time t :

$$n_i(t) = N_i(t) / \pi r^2, \quad (2)$$

where $N_i(t)$ represents the number of neighbors to particle i at time t . From this, we can get the fluctuation of the neighbor number density of particle i at time t .

$$\delta n_i(t) = \frac{n_i(t) - \bar{n}}{\bar{n}}, \quad (3)$$

where \bar{n} is the average number density of neighbors of N particles at M time points:

$$\bar{n} = \frac{1}{MN} \sum_{t=1}^M \sum_{i=1}^N n_i(t). \quad (4)$$

Based on this, the state of particle i at time t can be described as:

$$s_i(t) = \begin{bmatrix} \cos \theta_i(t) \\ \sin \theta_i(t) \\ \delta n_i(t) \end{bmatrix}, \quad (5)$$

where $\theta_i(t)$ is the movement direction angle of particle i at time t . Writing the state of N particles at time t together, and performing singular value decomposition after normalization:

$$A = \sum_{n=1}^r \sigma_n A_n^E = \sum_{n=1}^r \sigma_n \mathbf{v}_n \otimes \mathbf{u}_n, \quad (6)$$

$$A_{ij} = \sum_{n=1}^r \sigma_n (\mathbf{v}_n)_i (\mathbf{u}_n)_j. \quad (7)$$

Furthermore, the weight W_n^E of the eigen microstate in the Vicsek model can be obtained, where $W_n^E = \sigma_n^2$. In this way, the original Vicsek model can be used as an eigen microstate $U = [u_1, u_2, \dots, u_r]$ and the probability density distribution $w^E = [w_1^E, w_2^E, \dots, w_r^E]$ corresponding to each eigen microstate. Then, arranging the σ_n corresponding to the eigen microstates in order from large to small. The greater the value of σ_n , the greater the value of W_n^E .

3. Expansion of scaling analysis of eigen microstate method and group motion

In this chapter, we further analyze the phase transition of the Vicsek model under different noises, calculate the critical exponent of the Vicsek model, and study the phase transition of the Vicsek model under different densities.

3.1. Phase transition of Vicsek model under different noise

The probability amplitude σ_n of the eigen microstate u_n satisfies the scaling form:

$$\sigma_n(\eta, L) = L^{-\beta_\eta/\nu_\eta} f_l(h_\eta L^{1/\nu_\eta}), \quad (8)$$

Where L is the system scale, β_η is the critical exponent of σ_n under the influence of noise, ν_η is the critical exponent of the correlation length under the influence of noise, h_η is the distance from the noise to the phase transition point. The system weight factor satisfies the scaling form:

$$W_n^E(\eta, L) = \sigma_n^2(\eta, L) = L^{-2\beta_\eta/\nu_\eta} f_l^2(h_\eta L^{1/\nu_\eta}) = L^{-2\beta_\eta/\nu_\eta} F_l(h_\eta L^{1/\nu_\eta}). \quad (9)$$

Taking the partial derivative of h_η , then:

$$\frac{\partial W_n^E(\eta, L) L^{2\beta_\eta/\nu_\eta}}{\partial h_\eta} = L^{1/\nu_\eta} F_l'(h_\eta L^{1/\nu_\eta}). \quad (10)$$

The partial derivative of $W_n^E(\eta, L) L^{2\beta_\eta/\nu_\eta}$ with respect to h_η and the system scale L satisfy a power law relationship.




Figure 1. (Color online) *Log – Log* plot of W_l^E versus L around transition points: (a) W_1^E , $\eta_{1c} = 2.56$, $\beta_{1\eta}/\nu_{1\eta} = 0$; (b): W_2^E , $\eta_{2c} = 2.48$, $\beta_{2\eta}/\nu_{2\eta} = 0.94$; (c): W_3^E , $\eta_{3c} = \eta_{2c} = 2.48$, $\beta_{3\eta}/\nu_{3\eta} = \beta_{2\eta}/\nu_{2\eta} = 0.94$; (Li et al. [12].)

Taking the data of $\rho = 0.5$ as an example, as shown in Figure 1, the first largest eigen microstate of the Vicsek model undergoes a discontinuous phase transition at $\eta_{1c} = 2.56$. As shown in Figure 1, the second largest eigen microstate of the Vicsek model undergoes a continuous phase transition at $\eta_{2c} = 2.48$ and $\beta_{2\eta}/\nu_{2\eta} = 0.94$. As shown in Figure 1, the second largest eigen microstate of the Vicsek model undergoes a continuous phase transition at $\eta_{3c} = \eta_{2c} = 2.48$ and $\beta_{3\eta}/\nu_{3\eta} = \beta_{2\eta}/\nu_{2\eta} = 0.94$. It is consistent with the conclusion of the work of Li et al. [12]. As shown in Figure 2, the relationship between $W_2^E(\eta, L)L^{2\beta_\eta/\nu_\eta}$ and $h_\eta L^{1/\nu_\eta}$ is analyzed. It can be found that near the critical point, that is, within the range of $\eta \approx \eta_c$ and $h_\eta L^{1/\nu_\eta} \approx 0$, data at different scales can be scaled together. This means that within this range, the critical phase transition phenomenon of the Vicsek model still exists at different scales.




Figure 2. Log-Log plot of around transition points (a)The relationship between $W_2^E(\eta, L)L^{2\beta_\eta/\nu_\eta}$ and $h_\eta L^{1/\nu_\eta}$ in the second largest eigen microstate at $\rho = 0.5$; (b): Log – Log plot of the partial derivative of $W_2^E(\eta, L)L^{2\beta_\eta/\nu_\eta}$ with respect to h_η : $\frac{1}{\nu_\eta} = 1.20$. (Since the results in the simulation analysis are discontinuous, the difference is used instead of the partial derivative here.)

As shown in Figure 2, by calculating the partial derivative of $W_2^E(\eta, L)L^{2\beta_\eta/\nu_\eta}$ with respect to h_η , and analyzing it as the system scale L changes, it can be found that a power law relationship is satisfied between them. The reciprocal $1/\nu_n = 1.20$ of a straight line in Figure 2 whose slope is a critical exponent. The reciprocal of the critical exponent. Then the critical exponent $\beta_\eta = 0.783$ of the probability amplitude σ_n under the influence of noise and $\nu_\eta = 0.833$ of the correlation length under the influence of noise can be obtained.

4. Conclusion

This paper explores the scaling form and weight factor critical exponent of the continuous phase transition by an in-depth study of the microstate characteristics of the Vicsek model under different noise and density conditions, using the finite scale scaling method. The authors highlight the influence of density on the Vicsek model, revealing the phase transition behavior that emerges in different eigen microstates. By comparing the behavior under different parameter conditions, it can be found that noise play a key role in the Vicsek model.

References

- [1] Palacci J, Sacanna S, Steinberg A P, et al. Living Crystals of Light-Activated Colloidal Surfers [J]. *Science*, 2013, 339 (6122): 936–940.
- [2] Theurkauff I, Cottin-Bizonne C, Palacci J, et al. Dynamic Clustering in Active Colloidal Suspensions with Chemical Signaling [J]. *Physical Review Letters*, 2012, 108 (26): 268303.
- [3] Buttinoni I, Bialké J, Kümmel F, et al. Dynamical Clustering and Phase Separation in Suspensions of Self-Propelled Colloidal Particles [J]. *Physical Review Letters*, 2013, 110 (23): 238301.
- [4] Grégoire G, Chaté H. Onset of Collective and Cohesive Motion [J]. *Physical Review Letters*, 2004, 92 (2): 025702.
- [5] John T, Yuhai T. Long-Range Order in a Two-Dimensional Dynamical XY Model: How Birds Fly Together [J]. *Physical Review Letters*, 1995, 75 (23): 4326–4329.
- [6] Chaté H, Ginelli F, Grégoire G, et al. Modeling collective motion: variations on the Vicsek model [J]. *The European Physical Journal B*, 2008, 64 (3–4): 451–456.
- [7] Vicsek T, Czirok A, Ben-Jacob E, et al. Novel type of phase transition in a system of self-driven particles [J]. *Physical Review Letters*, 1995, 75 (6): 1226–1229.
- [8] Grégoire, G., Chaté, H. Onset of collective and cohesive motion [J]. *Physical Review Letters*, 2004, 92, 025702.
- [9] Chaté H, Ginelli F, Grégoire G, et al. Collective motion of self-propelled particles interacting without cohesion [J]. *Physical Review E*, 2008, 77, 046113.
- [10] Huepe C, Aldana M. New tools for characterizing swarming systems: A comparison of minimal models [J]. *Physica A*, 2008, 387, 2809–2822.
- [11] Wicks R T, Chapman S C, Dendy R O, Mutual information as a tool for identifying phase transitions in dynamical complex systems with limited data [J]. *Phys. Rev. E*, 2007, 75, 051125
- [12] Li X, Xue T, Sun Y, et al. Discontinuous and continuous transitions of collective behaviors in living systems [J], *Chinese Phys. B*, 2021, 30 128703