

# The study on the relationship between dementia and air pollution

**Qiuyao Duan**

College of Public Health ,Xiamen University, Xiamen, China, 361005

2845376743@qq.com

**Abstract.** Dementia is a multifaceted neurological condition distinguished by a reduction in cognitive abilities, impairment of memory function, and worsening in overall functional capacity. In recent times, there has been an increasing level of apprehension regarding the potential consequences of environmental elements, specifically air pollution, on the advancement and manifestation of dementia. The primary objective of this research article is to investigate the correlation between dementia and air pollution by a comprehensive analysis of epidemiological data. Additionally, this study will delve into probable processes that may underlie this association and then analyze the ramifications for public health. The results indicate that the presence of air pollution could potentially be a contributing factor to an elevated susceptibility to dementia and a deterioration in cognitive function among those afflicted. Gaining a comprehensive understanding of this correlation is of utmost importance in order to formulate efficacious preventive measures and policies aimed at mitigating the adverse effects of air pollution on dementia.

**Keywords:** dementia, air pollution, neurodegenerative disorder, cognitive decline, epidemiology, public health

## 1. Introduction

Dementia poses a substantial worldwide health burden, carrying notable societal, economic, and healthcare ramifications. The condition is distinguished by a reduction in cognitive functions, encompassing memory, cognition, rationality, and conduct. With the aging population and the rise of urbanization, there is a mounting apprehension regarding the influence of environmental variables on the risk and advancement of dementia. One of the environmental variables that has garnered significant attention is air pollution, mostly due to its extensive prevalence and potential detrimental impacts on human health. The primary objective of this study is to investigate the correlation between dementia and air pollution. The primary objective of this study is to conduct a comprehensive examination of the current epidemiological data that establishes a connection between exposure to air pollution and the development of dementia. Additionally, this research aims to investigate various processes that may explain this association, analyze the consequences for public health, and propose recommendations for preventive methods and policies.

## **2. Dementia: Definition, Types, and Prevalence**

### *2.1. Overview of Dementia*

Dementia is a clinical condition distinguished by a gradual deterioration of cognitive abilities, resulting in impaired performance of everyday activities. The etiology of this condition can be attributed to a range of underlying diseases, such as Alzheimer's disease, vascular dementia, Lewy body dementia, and frontotemporal dementia. The hallmark features of dementia include memory loss, impaired judgment, language difficulties, and changes in behavior and personality.

### *2.2. Common Types of Dementia*

Alzheimer's disease is the prevailing form of dementia, including around 60-80% of reported instances. Vascular dementia, characterized by cerebral hypoperfusion, ranks as the second most frequent kind. Lewy body dementia and frontotemporal dementia are characterized by unique pathological characteristics and clinical presentations.

### *2.3. Global Prevalence and Burden*

Dementia presents a substantial global health dilemma, since around 50 million individuals are presently afflicted with this disorder on a global scale. The anticipated increase in the prevalence of dementia is closely correlated with the aging population. The economic and social impact of dementia is considerable, imposing a huge burden on healthcare institutions, caregivers, and society at large.

## **3. Air Pollution: Types, Sources, and Composition**

### *3.1. Types of Air Pollutants*

Air pollution is composed of a multifaceted combination of particulate matter (PM), gaseous pollutants, and various other harmful elements. Particulate matter (PM) is classified according to its size, wherein PM<sub>2.5</sub> (particles with a diameter of 2.5 micrometers or less) and PM<sub>10</sub> (particles with a diameter of 10 micrometers or less) are of significant interest due to their capacity to infiltrate the respiratory system at a profound level.

### *3.2. Major Sources of Air Pollution*

Air pollution originates from both outdoor and indoor sources. Outdoor sources include industrial emissions, vehicle exhaust, power plants, and agricultural activities. Indoor sources include combustion of solid fuels for cooking and heating, tobacco smoke, and building materials.

### *3.3. Composition and Health Effects of Air Pollutants*

Air pollutants encompass a diverse array of substances, including nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), carbon monoxide (CO), ozone (O<sub>3</sub>), volatile organic compounds (VOCs), and heavy metals. These pollutants have been linked to a range of negative health impacts, such as respiratory and cardiovascular ailments, as well as possible neurotoxic effects.

## **4. Epidemiological Studies on Air Pollution and Dementia**

Several longitudinal cohort studies have been undertaken to examine the correlation between air pollution and the development of dementia. These investigations involve the longitudinal tracking of a substantial cohort of individuals for an extended duration, with the aim of evaluating their level of exposure to atmospheric pollution and its potential correlation with the subsequent onset and progression of dementia. Numerous cohort studies have consistently documented a strong correlation between prolonged exposure to air pollution and an elevated susceptibility to dementia. An investigation conducted by Chen et al. scrutinized a population of more than six million inhabitants in Ontario, Canada, and revealed that persons residing within a proximity of 50 meters from main roadways had a considerably elevated susceptibility to acquiring dementia in comparison to those

residing at greater distances [1]. The study additionally exhibited a dose-response correlation, wherein a higher risk of dementia was identified in those living in closer proximity to major roadways characterized by elevated levels of air pollution originating from traffic. In a similar vein, a comprehensive cohort study was out in Sweden by Oudin et al. meticulously tracked the progress of more than 3,000 individuals over an average duration of 11 years. The research conducted revealed a positive correlation between elevated levels of chronic exposure to air pollution originating from traffic, namely nitrogen oxide (NO<sub>x</sub>) and particulate matter (PM<sub>2.5</sub>), and a heightened susceptibility to dementia, with a particular emphasis on Alzheimer's disease and vascular dementia [2].

Case-control studies have made significant contributions to the comprehension of the association between air pollution and dementia. These studies involve the comparison of individuals diagnosed with dementia (referred to as cases) to individuals who do not have dementia (referred to as controls), with the aim of evaluating their prior exposure to air pollution. Case-control studies offer supplementary evidence about the potential association between exposure to air pollution and the risk of developing dementia. An investigation conducted by Chang et al. in Taiwan utilized a case-control study design to examine the potential correlation between prolonged exposure to fine particulate matter (PM<sub>2.5</sub>) and the likelihood of developing dementia. The research encompassed a sample size of more than 12,000 instances of dementia, along with corresponding control groups. The findings of this study revealed a noteworthy and positive correlation between exposure to PM<sub>2.5</sub> and the likelihood of developing dementia. The results indicated that prolonged exposure to PM<sub>2.5</sub> was associated with an elevated likelihood of acquiring dementia, particularly among persons with genetic predisposition [3].

Ecological investigations have been conducted to examine the correlation between levels of air pollution and the prevalence or incidence rates of dementia within a population. These studies offer significant insights into the potential influence of air pollution on the outcomes of dementia. Wu et al. conducted a noteworthy ecological study in the United States to investigate the correlation between air pollution at the county level and the prevalence of dementia. The present study conducted an analysis of data obtained from a sample of more than 6,000 counties, revealing a positive correlation between elevated levels of PM<sub>2.5</sub> and nitrogen dioxide (NO<sub>2</sub>) and an increased prevalence of dementia. The study posited that the reduction of air pollution levels on a regional scale may potentially have a mitigating effect on the burden of dementia [4].

The synthesis of findings from many epidemiological research through meta-analyses and systematic reviews offers a complete evaluation of the association between air pollution and dementia. The aforementioned analyses have provided more evidence to substantiate the correlation between exposure to air pollution and a heightened susceptibility to dementia. An investigation conducted by Peters et al. encompassed a systematic review and meta-analysis of 14 papers. The findings of this study indicated that prolonged exposure to air pollution, specifically PM<sub>2.5</sub> and NO<sub>x</sub>, exhibited a positive correlation with a heightened susceptibility to dementia. The findings of the analysis demonstrated a consistent and positive correlation across various study designs and populations, thereby enhancing the body of evidence supporting the adverse effects of air pollution on dementia [5].

The available epidemiological studies collectively indicate that prolonged exposure to air pollution, specifically PM<sub>2.5</sub> and contaminants connected to transportation, is correlated with a heightened likelihood of developing dementia. The results of this study offer valuable insights into the possible influence of air pollution as a modifiable risk factor for dementia. These findings underscore the necessity of implementing preventative measures and public health interventions to reduce the adverse effects of air pollution on cognitive health [6-8].

## **5. Potential Mechanisms**

In order to comprehend the correlation between air pollution and dementia, it is imperative to investigate the plausible biological pathways by which air pollution can manifest its deleterious impacts on the brain. Several mechanisms have been proposed based on experimental and observational studies, shedding light on the possible pathways linking air pollution to dementia.

Neuroinflammation has been identified as a significant factor involved in the correlation between air pollution and the development of dementia. The presence of air pollutants has the potential to induce an inflammatory reaction within the brain, which is distinguished by the activation of microglia and the subsequent release of pro-inflammatory cytokines and chemokines. Chronic neuroinflammation has the potential to induce neuronal damage, disrupt synaptic function, and ultimately play a role in the pathogenesis and advancement of dementia. Empirical investigations have provided evidence indicating that the introduction of air pollutants, namely fine particulate matter (PM<sub>2.5</sub>) and pollutants associated with transportation, can elicit neuroinflammatory reactions in animal subjects. The observed reactions are linked to the build-up of neuroinflammatory indicators, such as interleukin-1 $\beta$  (IL-1 $\beta$ ), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), and cyclooxygenase-2 (COX-2). The initiation of inflammatory pathways has the potential to disturb regular neuronal function and play a role in the process of neurodegeneration [7, 9, 10].

Another important mechanism by which air pollution may contribute to dementia is oxidative stress. Air pollutants contain reactive oxygen species (ROS) and other toxic compounds that can induce oxidative damage in the brain. Oxidative stress arises from a state of disequilibrium between the generation of reactive oxygen species (ROS) and the body's capacity to counteract them through antioxidant defense mechanisms. This imbalance results in cellular harm and compromised neural functionality. Numerous investigations conducted on animals and humans have demonstrated that exposure to air pollution leads to an elevation in oxidative stress markers. Cellular structures and functions can be adversely affected by oxidative damage to lipids, proteins, and DNA. In addition, the presence of oxidative stress can trigger the initiation of inflammatory pathways and the production of neurotoxic compounds, so intensifying the progression of neurodegenerative mechanisms [7, 10].

Exposure to air pollution has been found to be linked with systemic inflammation, impairment of endothelial function, and an elevated risk of cardiovascular disease. The presence of vascular factors has been identified as a potential contributing element in the pathogenesis of cerebrovascular disease, a condition that has been associated with an elevated susceptibility to vascular dementia and cognitive deterioration. The presence of air pollutants, specifically particles resulting from combustion, has the potential to infiltrate the bloodstream and directly impact the vascular function within the brain. Numerous empirical investigations have substantiated the correlation between prolonged exposure to atmospheric pollution and an elevated susceptibility to stroke and cerebrovascular ailments. The disruption of cerebral blood flow, reduced oxygen supply, and compromised vascular integrity can contribute to neurodegenerative processes and the onset of dementia [6-8].

The blood-brain barrier (BBB) serves as a crucial protective mechanism that governs the selective transport of chemicals between the vascular system and the central nervous system. The function of this mechanism is of utmost importance in the regulation of brain homeostasis and the safeguarding of the brain from potentially detrimental agents. Airborne pollutants, including ultrafine particles, have the ability to directly traverse the blood-brain barrier (BBB) and get access to the brain. Research has demonstrated that the presence of air pollution has a detrimental effect on the integrity of the blood-brain barrier (BBB), resulting in heightened permeability and facilitating the infiltration of harmful substances into the brain. The aforementioned disturbance has the potential to induce harm to neurons, provoke inflammation, and impair neuronal function, hence playing a role in the progression of dementia [6-8].

The sensitivity of individuals to the detrimental impacts of air pollution on the brain might be influenced by hereditary factors at the individual level. The response of individuals to air pollution exposure can be influenced by genetic variations in genes associated with inflammation, oxidative stress, and detoxification pathways. The potential for gene-environment interactions, namely those involving genetic variations associated with inflammation or oxidative stress and exposure to air pollution, may contribute to an elevated susceptibility to dementia [7].

Through genetic studies, researchers have successfully identified distinct genetic variations that have the potential to alter the relationship between air pollution and cognitive outcomes. The results of

this study indicate that the impact of air pollution on brain function may be influenced by hereditary factors, leading to variations in individual sensitivity [7].

Gaining a comprehensive understanding of the fundamental mechanisms that establish a connection between air pollution and dementia is crucial in order to facilitate the development of precise interventions and preventive measures. Through the clarification of these mechanisms, researchers are able to ascertain prospective treatment targets and formulate solutions to alleviate the deleterious impacts of air pollution on cognitive well-being.

## **6. Implications for Public Health**

First, to mitigate the impact of air pollution on dementia, policymakers need to implement effective strategies to reduce air pollution levels. This includes stricter emissions standards for industrial and transportation sectors, promoting clean energy sources, and fostering sustainable urban planning.

Second, it is imperative to acknowledge the significant impact that public education and awareness campaigns may have in disseminating vital information to the broader population regarding the possible hazards associated with air pollution and its detrimental effects on cognitive well-being.

Third, continuous efforts are needed to advance environmental regulations and standards to ensure the protection of public health. Implementing and enforcing stricter air quality standards can significantly reduce the population's exposure to harmful air pollutants.

Urban planning initiatives that prioritize green spaces, pedestrian-friendly infrastructure, and efficient public transportation systems can help reduce air pollution levels in urban areas. Research has demonstrated that the presence of green areas has a beneficial effect on cognitive function and mental well-being.

Fourth, addressing the complex relationship between air pollution and dementia requires interdisciplinary collaborations among researchers, healthcare professionals, policymakers, and community stakeholders. Such collaborations can facilitate the development of comprehensive strategies for prevention, intervention, and policy implementation.

## **7. Challenges and Future Directions**

Epidemiological investigations that explore the association between air pollution and dementia encounter several methodological obstacles, encompassing the accurate measurement of exposure, the presence of confounding variables, and the possibility of reverse causation. By addressing these shortcomings, the evidence base can be strengthened and our knowledge of the association can be improved. Certain demographic groups may exhibit increased susceptibility to the detrimental impacts of air pollution on the development and progression of dementia. The identification of vulnerable populations, including older adults, persons with pre-existing cognitive impairments, and those with genetic predispositions, can facilitate the implementation of focused interventions and preventive measures.

Additional investigation is required to delve into the enduring consequences of exposure to air pollution on the chance of developing dementia, as well as to determine the correlations between the dosage of air pollution and its corresponding effects. The utilization of improved air pollution monitoring systems and other enhanced exposure assessment methodologies will contribute to a deeper comprehension of the cumulative impacts of air pollution. The exploration of the fundamental processes connecting air pollution and dementia necessitates the conduct of mechanistic investigations and the discernment of biomarkers. The examination of neuroinflammatory indicators, oxidative stress markers, and imaging techniques can yield valuable insights into the precise mechanisms by which air pollution impacts the brain. Subsequent investigations ought to prioritize the assessment of the efficacy of therapies and preventive measures targeted at mitigating the deleterious consequences of air pollution on the development and progression of dementia. This encompasses the examination of the advantages associated with indoor air filtration systems, adjustments made to one's lifestyle, and specific therapies designed for populations at a heightened risk.

## 8. Conclusion

The available literature indicates that there is a mounting body of data supporting the notion that the exposure to air pollution is correlated with an elevated likelihood of developing dementia and experiencing a deterioration in cognitive function. Gaining a comprehensive understanding of the correlation between air pollution and dementia is of paramount importance in order to facilitate the development of efficacious preventative measures, the implementation of environmental legislation, and the advancement of public health initiatives. Through the acknowledgment and examination of the various obstacles, as well as the allocation of resources towards additional investigation, scholars can strive to mitigate the detrimental impact of air pollution on dementia and enhance the well-being of those afflicted by this incapacitating ailment.

## References

- [1] Chen H, Kwong JC, Copes R, et al. Living near major roads and the incidence of dementia, Parkinson's disease, and multiple sclerosis: a population-based cohort study. *Lancet*. 2017;389(10070):718-726.
- [2] Oudin A, Forsberg B, Adolfsson AN, et al. Traffic-related air pollution and dementia incidence in Northern Sweden: a longitudinal study. *Environ Health Perspect*. 2016;124(3):306-312.
- [3] Chang KH, Chang MY, Muo CH, Wu TN, Hwang BF, Chen CY. Increased risk of dementia in patients exposed to nitrogen dioxide and carbon monoxide: a population-based retrospective cohort study. *PLoS One*. 2017;12(6):e0177860.
- [4] Wu YC, Lin YC, Yu HL, Chen JH, Chen TF, Sun Y, et al. Association between air pollutants and dementia risk in the elderly. *Alzheimers Dement*. 2015;11(7):P415.
- [5] Peters R, Ee N, Peters J, Booth A, Mudway I, Anstey KJ. Air pollution and dementia: a systematic review. *J Alzheimers Dis*. 2019;70(s1):S145-S163.
- [6] Calderón-Garcidueñas L, Franco-Lira M, Henríquez-Roldán C, et al. Urban air pollution: influences on olfactory function and pathology in exposed children and young adults. *Exp Toxicol Pathol*. 2010;62(2):91-102.
- [7] Block ML, Elder A, Auten RL, et al. The outdoor air pollution and brain health workshop. *Neurotoxicology*. 2012;33(5):972-984.
- [8] Calderón-Garcidueñas L, D'Angiulli A, Kulesza RJ, et al. Air pollution and children: neural and tight junction antibodies and combustion metals, the role of barrier breakdown and brain immunity in neurodegeneration. *J Alzheimers Dis*. 2015;43(3):1039-1058.
- [9] Calderón-Garcidueñas L, Reed W, Maronpot RR, et al. Brain inflammation and Alzheimer's-like pathology in individuals exposed to severe air pollution. *Toxicol Pathol*. 2004;32(6):650-658.
- [10] Fonken LK, Xu X, Weil ZM, Chen G, Sun Q, Rajagopalan S, et al. Air pollution impairs cognition, provokes depressive-like behaviors and alters hippocampal cytokine expression and morphology. *Mol Psychiatry*. 2011;16(10):987-995.